FUNGICIDES FOR MUSHROOM CROPS Apr-18							
Active Ingredient	Target Organism	Production area m2	Registration/ Status	Comments			
prochloraz managnese	Wet Bubble, Dry Bubble, Cobweb	on casing 17 500m2/wk	registered in SA	Frequently used by most growers; In Europe supplied by BASF			
prochloraz zinc	Wet Bubble, Dry Bubble, Cobweb	on casing 17 500m2/wk	registered in SA	Used by many growers, cheaper than Octave/Sporgon			
thiabendazole	Wet Bubble, Dry Bubble, Cobweb	on casing 17 500m2/wk	registered in SA				
metafenone	Cobweb, Dry Bubble	on casing 17 500m2/wk	not registered	Registered in Europe; used in vine yards			
carbendazim	Trichoderma, Cobweb, Dry- & Wet Bubble	on casing 17 500m2/wk	not registered	Carbendazim is used on beds as casing treatment (EC or SC); the Australian dept Agric has already permitted the use of Carbendazim on agaricus mushrooms, and the toxicology data is available.			
benomyl	Trichoderma spp		not registered	Used as a spawn (seed) treatment at spawning and in small quantities thus giving very low residue levels.			
Safe Tray, Bumber, Banner, Propicon	Mushroom mycelium transferring virus	16% of growers use trays	not registered	To dip wooden trays to protect against various fungi and mycelium transferring virus and; used in Australia			

BIO-CONTROL PRODUCTS

IBacillus amyloliquefaciens	Trichoderma compost mould in phase II		not registered	registered and used in States
-----------------------------	--	--	----------------	-------------------------------